

Horticultural interventions may reduce adults' depressive symptoms: A systematic review of randomized controlled trials

Claudio D. Rosa^{a,*}, Talisson S. Chaves^b, Silvia Collado^c, Lincoln R. Larson^d, KangJae Jerry Lee^e, Christiana C. Profice^a

^a Department of Development and Environment, State University of Santa Cruz, Brazil

^b Department of Physiological Sciences, Federal University of São Carlos, Brazil

^c Department of Psychology and Sociology, University of Saragossa, Spain

^d Department of Parks, Recreation and Tourism Management, North Carolina State University, United States

^e Department of Parks, Recreation and Tourism, University of Utah, United States

ARTICLE INFO

Handling editor: L. McCunn

Keywords:

Depressed
Gardening
Horticulture
Mental health

ABSTRACT

We conducted a systematic review to examine the effect of horticultural interventions (e.g., planting or taking care of plants) on people's depressive symptoms as assessed by depression outcome measures. On January 19 of 2022, the databases MEDLINE (PubMed), PsycArticles (APA), SCOPUS (Elsevier), Google Scholar, and ClinicalTrials.gov were searched from inception. The decision to include or exclude studies in the full text, the data extraction, and the risk of bias assessment were performed by two researchers. We identified 20 randomized controlled trials (RCTs) ($n = 998$ participants; all adults), from nine different countries. Overall, we found evidence that some horticultural interventions plus usual care (i.e., continuing normal routine for healthy people or treatment for unhealthy ones) may reduce depressive symptoms more than usual care alone, with most studies suggesting a moderate (Hedges' $g \geq 0.5$) or large effect ($g \geq 0.8$). The percentage of participants who dropped out from any of the horticultural interventions ranged from 0% to 40% and only one study reported adverse events (i.e., fatigue and tiredness) related to the intervention. Except for one study, all studies had some risk of bias due to design limitations, such as lack of participants' blinding and/or a prespecified analysis plan. Our findings suggest that some horticultural interventions are effective and safe to use as a complementary strategy to reduce adults' depressive symptoms. More RCTs are needed to understand how specific participants and intervention characteristics can alter the effect of horticultural interventions on depressive symptoms.

1. Introduction

Depression is one of the most serious global health challenges (Cipriani et al., 2018). Before the COVID-19 pandemic, it was estimated that 322 million people in the world dealt with this disorder, which can harm different dimensions of people's lives including affective relationships, professional achievement, and overall health and well-being (World Health Organization, 2017). Unfortunately, this prevalence may now be higher since a 27.6% increase in depression was associated with the COVID-19 pandemic (World Health Organization, 2022). The use of antidepressants and psychotherapy are two of the most well-known and recommended treatments for depression (Lopresti, 2019). Nonetheless, even the combination of these treatments commonly produces small improvements in depressive symptoms (Cuijpers et al., 2020; Lopresti,

2019; McCormack & Korownyk, 2018). Thus, efforts have been directed towards complementary interventions that may help to provide greater reductions in depressive symptoms, such as physical exercise (Catalan-Matamoros et al., 2016), diet changes (Berk & Jacka, 2019), and contact with nature (Rosa et al., 2021). The use of nature-based activities to reduce people's depressive symptoms seems especially promising when compared to physical exercise and diet changes (Rosa et al., 2021). For example, Rosa et al. (2021) found that, compared to usual care, participants in forest therapy groups were 17 times as likely to achieve remission and three times as likely to have at least a 50% reduction on depressive symptoms.

Several theories and frameworks have been used to explain the health benefits associated with activities in nature (Fernee et al., 2017; Houge Mackenzie et al., 2021; Kaplan, 1995; Reese & Gosling, 2020;

* Corresponding author. Department of Development and Environment, State University of Santa Cruz, Ilhéus, Bahia, Brazil.

E-mail address: claudio2008ilheus@hotmail.com (C.D. Rosa).

<https://doi.org/10.1016/j.jenvp.2023.102112>

Received 28 February 2023; Received in revised form 11 July 2023; Accepted 14 August 2023

Available online 18 August 2023

0272-4944/© 2023 Elsevier Ltd. All rights reserved.

Russell & Farnum, 2004; Ulrich et al., 1991; Wilson, 1984). Among these, Attention Restoration Theory (ART, Kaplan, 1995) and Stress Recovery Theory (SRT, Ulrich et al., 1991) have emerged as the most popular theoretical explanations (Berto, 2014; Crossan & Salmoni, 2021; Frost et al., 2022; Hartig, 2021; Jiang et al., 2021; Moll et al., 2022; Ohly et al., 2016). Taken together ART and SRT posit that positive experiences in nature can be pleasurable, reduce anxiety and stress, and improve concentration and mood, all of which are related to lower depressive symptomatology (Fried, 2017; Kaplan, 1995; Owens & Bunce, 2022; Rosa et al., 2021; Ulrich et al., 1991). In accordance with these theories, research suggests that some activities involving contact with nature may improve people's depressive symptoms such as sad mood (Soga et al., 2017), difficulty in concentrating (Clatworthy et al., 2013), sleep problems (Shin et al., 2012), and hopelessness (Sturm et al., 2012). Despite the potential benefits of nature-based activities, systematic reviews on the effect of nature-based interventions on depression are scarce, hindering our knowledge about what types of nature-based activities (if any) are best to improve depressive symptoms.

Three different types of nature-based interventions are often described in the academic literature: forest therapy (e.g., Kim et al., 2009), nature-based adventure (e.g., Sturm et al., 2012), and horticultural activities (e.g., Kam & Siu, 2010). Systematic reviews were already done to investigate the effect of the first two types of nature-based interventions on depression (Rosa, Chaves, Collado, Larson, & Profice, 2023; Rosa et al., 2021) but, to our knowledge, the effect of horticultural activities on depressive symptoms has not been systematically reviewed. We use horticultural interventions as a broad term encompassing both horticultural therapy and therapeutic horticulture. According to the American Horticultural Therapy Association (AHTA, 2017, p.2), "horticultural therapy is the participation in horticultural activities facilitated by a registered horticultural therapist to achieve specific goals within an established treatment, rehabilitation, or vocational plan" while therapeutic horticulture is "the participation in horticultural activities facilitated by a registered horticultural therapist or other professionals with training in the use of horticulture as a therapeutic modality to support program goals". Thus, we use the term horticultural intervention to refer to any horticultural activity facilitated by a horticultural therapist or other trained professional to achieve health benefits (AHTA, 2017). Examples of horticultural interventions include planting and taking care of plants with the support of a therapist or other trained professional (Soga et al., 2017).

Although many reviews have assessed the effect of horticulture on health-related outcomes (Cipriani et al., 2017; Clatworthy et al., 2013; Kamioka et al., 2014; Liu et al., 2014; Murrioni et al., 2021; Nicholas et al., 2019; Soga et al., 2017; Tu, 2022; Wang & MacMillan, 2013; Wang et al., 2022), no studies have conducted a systematic review of the effect of horticultural interventions on people's depressive symptoms. For instance, Nicholas et al. (2019) conducted a systematic review to assess the effect of horticultural therapy on older adults, but their review identified only four primary studies evaluating the effect of horticultural therapy on depressive symptoms as assessed by depression outcome measures. This small number of identified studies can be partially explained by the authors' eligibility criteria that excluded studies with younger adults, adolescents, and children, studies not published in English, and those published before January 2008. Importantly, this small pool of research (i.e., only four studies) constitutes a fraction of the existing empirical evidence on the effect of horticultural interventions on people's depressive symptoms. The lack of a systematic synthesis of previous research hinders practitioners to develop guidelines and effective intervention programs that can prevent or treat depression (Owens & Bunce, 2022; Rosa et al., 2021). We therefore present a systematic review summarizing the effect of horticultural interventions on depressive symptoms. Our broad eligibility criteria (e.g., including studies in any language and from any period of time), together with a search strategy focused on depression, allowed us to identify more

studies assessing the effect of horticultural interventions on depression than any previous systematic review. We also collected information about dropouts and adverse events. Our systematic synthesis is expected to deepen the understanding of the potential utility of horticultural interventions in reducing depressive symptoms. The overarching research question guiding our review was: "What is the effect of horticultural interventions on depressive symptoms as compared to alternative interventions (or no intervention)?"

2. Method

2.1. Eligibility criteria

The criteria for inclusion in our review are summarized in Table 1, and a detailed description of these criteria can be found in our registered protocol (Supplementary File 1), which was built based on the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) 2015 statement (Shamseer et al., 2015). We did not exclude studies based on language, date, or because they were not published in a peer-reviewed journal. Although the aim of our study was not restricted to adults, we were only able to identify eligible randomized controlled trials (RCTs) conducted with this age group.

In this study, we focus on RCTs. We did this because randomization ensures that any differences between groups in prognostic/confounding variables at the baseline are due to chance (Sterne et al., 2016, 2019).

2.2. Search strategy

We used previous systematic reviews on related topics (e.g., the effects of horticultural therapy on older adults' health) as an informative source to identify eligible primary studies (e.g., Murrioni et al., 2021; Nicholas et al., 2019), and we searched for primary studies that were not included in these systematic reviews. On January 19 of 2022 the databases MEDLINE (PubMed), PsycArticles (APA), SCOPUS (Elsevier), Google Scholar, and ClinicalTrials.gov were searched from inception. Additionally, we checked the references of included studies and our personal files (e.g., computer archives), which could provide access to additional studies. Our exact search strategy is described in our registered protocol (Supplementary File 1).

2.3. Selection, data extraction, and risk of bias assessment

The first author performed the title and abstract screening, selection based on full-text, data extraction, and risk of bias assessment. Another researcher checked whether the eligibility criteria were applied appropriately, and also examined the data extraction and the risk of bias

Table 1

Eligibility criteria for our review based on population (P), intervention (I), comparison groups of interest (C), outcomes (O), and study designs (S).

PICOS	Description
Population	Studies with humans at any age, healthy or unhealthy
Intervention	Any horticultural activity facilitated by a horticultural therapist or other trained professional to achieve health benefits.
Comparison groups of interest	Studies with any comparison/control group and studies without a control group.
Outcomes	Studies that assess depression using a measure designed to measure depression. At least one study (i.e., a validation study) should exist describing how the content of the measure matches the construct's content (i.e., depression).
Study design ^a	Randomized and non-randomized studies of interventions.

Note.

^a It was part of our eligibility criteria to include both randomized and non-randomized studies of interventions. In this manuscript, we focus on randomized controlled trials (RCTs). The findings from non-randomized studies will be reported in a separate study.

assessment. Specifically, the second researcher read through the decisions made by the first researcher and approved/disapproved them. The few disagreements between the first author and the other researcher were resolved through discussion. From each study, we collected information regarding participants' sociodemographic variables, the setting where the interventions took place, the horticultural activities conducted, and the depression score at baseline and after the intervention (see Table 2 in Supplementary File 1). The studies' risk of bias was assessed with the RoB 2 tool (Sterne et al., 2019).

2.4. Data synthesis

To estimate the effect of horticultural interventions on depressive symptoms, we extracted data from the pre-test closest to the start of the intervention and the post-test closest to the end of the intervention. When studies used more than one depression outcome measure, we selected just one measure based on pre-specified criteria (see "Dealing with Multiple Effect Estimates" in Supplementary File 1). When possible, we calculated Hedges' *g* using each group's mean change in depression scores from pre to post-intervention and its standard deviation. Otherwise, we calculated *g* by using the post-test scores and its standard deviation (Higgins et al., 2019). Although depression outcome measures varied, we were able to calculate the percentage of change in depressive symptoms from baseline to post-intervention and the standardized mean change (as described by Morris, 2008) in the majority of studies. Focusing on change in depressive symptoms is more appropriate than the difference between groups following treatment when group scores differ substantially at baseline (Vickers, 2001).

Another relevant outcome was the number of participants who demonstrated substantial improvement following the intervention. We operationalized response to the intervention as a $\geq 50\%$ decrease in depressive symptoms from baseline (Riedel et al., 2010). Research shows that a $\geq 50\%$ decrease is a good proxy for clinically relevant improvement in depression as assessed by three depression scales: Hamilton Depression Rating Scale (HDRS), Beck Depression Inventory (BDI), and Montgomery Asberg Depression Rating Scale (MADRS) (Riedel et al., 2010). Accordingly, we calculated the number of participants reporting a $\geq 50\%$ decrease in depressive symptoms when the studies used one of these three scales. This number was calculated using the formula described by Furukawa et al. (2005). For all studies with available data, we report the number of participants who dropped out and the adverse events that occurred. When feasible, we calculated risk ratios for dichotomous outcomes because these are easier to understand than odds ratios (Higgins et al., 2019).

We conducted a fixed-effects meta-analysis to avoid an over-estimation of the intervention effect due to a huge effect observed by one study, and we ran sensitivity analyses to check the robustness of our findings (Higgins et al., 2019). In this meta-analysis, we also assessed whether the results from studies that offered other interventions (co-interventions) in addition to horticulture revealed greater improvements in people's depressive symptoms than studies that just involved horticulture.

Because no study reported having substituted participants' usual treatment with horticultural activities, we assumed that the horticultural interventions were used as a complementary intervention for unhealthy participants or as the only intervention for healthy ones. We used the term "usual care" to represent individuals' keeping their normal routine; this normal routine means that unhealthy participants continued their usual treatment (e.g., psychotherapy), and the healthy ones received no intervention. To clarify the distinction between usual care and co-interventions, we use Kim et al.'s (2016) study as an example. This study was conducted with patients with Alzheimer at Seongdong-gu Center for Dementia. Usual care in this case is the normal care offered to patients at this center and co-interventions are the additional interventions (e.g., exercise and music therapy), other than horticulture, provided to the study's participants.

To facilitate the interpretation of the findings from the RCTs included in this systematic review, we report estimates of effects and, when feasible, 95% confidence intervals (CI) for these estimates. Hedges' *g* and risk ratios were calculated using RevMan (Review Manager (RevMan) [Computer Program], 2020), and figures illustrating the risk of bias of RCTs were created using robvis (McGuinness, 2019). All data utilized in our analyses that are not reported in the manuscript are available in Supplementary File 2. This file also contains the references for all randomized studies included in our systematic review.

3. Results

Our database searches produced 223 records, from which 62 were deemed eligible after the full-text assessment. An example of a study excluded after the full-text assessment is Shao et al. (2020), who did not assess people's depression using a depression outcome measure. An additional 20 studies were identified through supplementary search strategies such as checking the reference list of all eligible studies and previous systematic reviews on related topics (e.g., Nicholas et al., 2019; Soga et al., 2017). Thus, a total of 82 studies were deemed eligible based on our eligibility criteria (Table 1). From these 82 eligible studies, 20 were RCTs that were considered in the present study (see Fig. 1 for a flow diagram). These 20 RCTs took place in nine different countries and involved a total of 998 adults (Table 2). Sixteen studies were conducted in Asia, three in Europe, and one in the United States of America. All studies were published in peer-reviewed scientific journals from 2003 to 2021, with more than half published in the last five years (2017–2021). These studies included older and middle-aged adults, psychiatric and stroke patients, and university students. No study included children or adolescents.

Horticultural interventions involved a variety of activities, such as sowing, potting, planting, making bouquets, making a terrarium, watering plants, and harvesting (Table 3). Also variable was the length, frequency, and duration of these interventions. Intervention length varied from two to 26 weeks. The frequency of horticultural interventions ranged from weekly to daily sessions, and duration from one to 4 h. Some horticultural interventions were associated with co-interventions such as physical activities, cognitive occupational therapy, art therapy, stress management lessons, and physiotherapy. The effects of horticultural interventions were most often compared with usual care but they were also compared with other interventions like educational sessions, exercise therapy, social activities, other occupational activities, and stress management sessions. Seven different measures were used to assess depression. The short form of the Geriatric Depression Scale was the one most frequently used (Table 2). We were able to evaluate the risk of bias of 19 RCTs, from which we deemed 18 as at a "high" risk of bias and one at "low" risk of bias (Fig. 2). One study was not evaluated because we only had access to its abstract (Moshfeghi et al., 2014).

3.1. Horticultural intervention versus usual care alone

Overall, 15 RCTs compared horticultural interventions plus usual care with usual care only. Of the 15 RCTs providing data for this comparison, 13 suggest that horticultural interventions plus usual care may reduce depressive symptoms more than usual care alone, including 12 studies that provided data for a fixed-effects meta-analysis (Hedges' *g* = -1.26 , 95% CI [-1.47 , -1.05], $p < .001$, $I^2 = 91.9\%$). Eleven of these 12 studies reported a moderate ($g \geq 0.5$) or large ($g \geq 0.8$) effect size (Fig. 3a).

Studies in which participants took part in horticultural interventions plus additional interventions (i.e., co-interventions) like physiotherapy resulted in a smaller combined estimate than the one obtained from studies in which a horticultural intervention was the only reported intervention (Fig. 3a). We conducted two sensitivity analyses to understand the robustness of these findings. First, we ran a random-effects




Fig. 1. Flowchart illustrating the process of identifying and selecting studies.

meta-analysis to check the impact of statistical heterogeneity in our results. This analysis produced similar results to the ones observed in Fig. 3 with an even larger combined estimate because the random effects meta-analysis gave more weight to a single study that found a very large estimate of effect (Chu et al., 2019). Second, after removing this single study from the meta-analysis, the effect remained large and in the same direction. In other words, regardless of method, we observed a large effect favoring horticultural interventions, suggesting these findings are robust.

Only three studies reported data necessary to estimate the number of participants who had a $\geq 50\%$ reduction in their depressive symptoms from baseline to post-intervention (Fig. 3b). The combined estimate from a fixed-effects meta-analysis of these studies suggests that participants in the horticultural interventions were twice as likely to have a $\geq 50\%$ reduction in their depressive symptoms from baseline to post-intervention than participants only receiving usual care (Risk Ratio = 2.03 [1.38, 2.98], $p = .002$, $I^2 = 84\%$). Similar to the previous meta-analysis (Fig. 3a), we ran additional tests to check the robustness of these findings. First, a random-effects meta-analysis suggested an even larger estimate, but the 95% CI was much larger due to statistical heterogeneity (Risk Ratio = 2.77, [0.36, 21.03], $p = .32$, $I^2 = 84\%$). Second, after removing a study that found a very large estimate of effect

(Ghanbari et al., 2015) from these meta-analyses, the combined estimate of effect became smaller and statistical heterogeneity disappeared (Risk Ratio = 1.28 [0.93, 1.72], $p = .11$, $I^2 = 0$), suggesting these findings are not robust.

Among the RCT studies that compared horticultural interventions plus usual care to usual care only but did not report data to be included in the meta-analysis, Moshfeghi et al. (2014) reported that their horticultural intervention group had a statistically significant larger reduction in the mean depression score than their control group ($p < .01$). In addition, two studies found non-statistically significant differences in depressive symptoms between the horticultural interventions and the usual care groups (Ng et al., 2018; Pálsdóttir et al., 2020).

Ten studies reported the number of participants who dropped out from horticultural interventions and the number of participants who dropped out from the usual care groups (Table 4). In eight studies, no dropout occurred. The two studies that reported dropouts pointed in opposite directions: one study found that more participants dropped out from the horticultural intervention group (Risk Ratio = 5.00 [0.27, 94.34], $p = .28$), and the other found that fewer participants dropped out from the horticultural intervention group (Risk Ratio = 0.14 [0.02, 1.10], $p = .06$). Overall, dropouts from horticultural interventions ranged from zero to 40%. Only one study reported an adverse event

Table 2

Main characteristics of the randomized controlled trials (RCTs) included in this systematic review of studies investigating the effects of horticultural interventions on depressive symptoms.

First author (year)	Participants	Mean age or age range	Women %	Time (T) in which data was collected ^b	Depression measure	Country	Setting where the horticultural intervention took place
Buru et al. (2021)	University students	20.2	Unclear	T1: Before the intervention T2: After the intervention	Beck Depression Inventory	Romania	University of Agricultural Sciences and Veterinary Medicine
Pálsdóttir et al. (2020)	Stroke survivors	67	60	T1: Before the intervention T2: Eight months after randomization	Hospital Anxiety and Depression Scale	Sweden	Alnarp Rehabilitation Garden
Kim et al. (2020a)	Elderly living in a homeless living facility	73.2	33.3	T1: Before the intervention T2: After the intervention	Geriatric Depression Scale-Short Form	South Korea	In some parts of the garden
Kim et al. (2020b)	Caregivers of elderly with dementia	60.0	100	T1: Before the intervention T2: After the intervention	Center for Epidemiological Studies Depression Scale	South Korea	At a health center
Makizako et al. (2020)	Older adults with depressive symptoms and memory problems	73.1	50.6	T1: Before the intervention T2: Immediately after the intervention	Geriatric Depression Scale-Short Form	Japan	Public garden
Chu et al. (2019)	Older residents of nursing homes	78.6	62.7	T1: Before the intervention T2: At the end of the intervention	Geriatric Depression Scale-Short Form	China	Indoors at a table where residents could sit
Najjar et al. (2018)	Chronic depressed male outpatients	Unclear	0.0	T1: Before the intervention T2: After the intervention	Depression, Anxiety, and Stress Scale-44	Iran	Noor-Almahdi Mental Hospital
Kim (2018)	Middle-aged women	40 to 59	100	T1: Before the intervention T2: After the intervention	Zung Self-rating Depression Scale	South Korea	At a culture center in Incheon
Lai et al. (2018)	Frail and prefrail nursing home residents	84.6	65.6	T1: Before the intervention T2: Immediately after the intervention	Geriatric Depression Scale-Short Form	China	Indoor and outdoors
Ng et al. (2018)	Older adults	67.7	78.0	T1: At the start of the study T2: Three months after the intervention	Zung Self-rating Depression Scale	Singapore	Indoor and outdoor activities at parks, gardens, and a nature reserve
Vujcic et al. (2017)	Psychiatric patients	45.4	70.0	T1: Before the intervention T2: Directly after the intervention	Depression Anxiety Stress Scale-21	Serbia	The Jevremovac Botanical Garden
Kim et al. (2016)	Patients with Alzheimer	78.5	69.8	T1: Before the intervention T2: After the intervention	Geriatric Depression Scale-Short Form	South Korea	Seongdong-gu Center for Dementia
Detweiler et al. (2015)	War veterans	46.4	4.2	T1: Before the intervention T2: After the intervention	Center for Epidemiological Studies Depression Scale	United States of America	Veterans Affairs Medical Center in Salem, Virginia
Ghanbari et al. (2015)	Female students of Golestan dormitory	20.6	100	T1: Before the intervention T2: After the intervention	Beck Depression Inventory	Iran	Dormitory yard
Kotozaki et al. (2015)	Women victims of an earthquake	43.4	100	T1: Before the intervention T2: After the intervention	Center for Epidemiologic Studies Depression Scale	Japan	At a university lab and at participants' homes
Kotozaki (2014)	Women victims of an earthquake	46.5	100	T1: First day of the intervention T2: After the intervention	Center for Epidemiologic Studies Depression Scale	Japan	At a community center and at participants' homes
Moshfeghi et al. (2014) ^a	Older adults in nursing homes	Unclear	Unclear	T1: Before the intervention T2: After the intervention	Depression, Anxiety, and Stress Scale-44	Iran	Unclear
Tse and Ho (2013)	Older persons living in nursing homes	60 to 89	62.2	T1: Before the intervention	Geriatric Depression Scale-Short Form	China	Nursing home

(continued on next page)

Table 2 (continued)

First author (year)	Participants	Mean age or age range	Women %	Time (T) in which data was collected ^b	Depression measure	Country	Setting where the horticultural intervention took place
Kam and Siu (2010)	People with psychiatric illness	44.3	29.7	T1: Before the intervention T2: After the intervention	Depression Anxiety Stress Scale-21	China	New Life Farm
Kim et al. (2003)	Poststroke hemiplegic patients	56.0	30.95	T1: Before the intervention T2: After the intervention	Beck Depression Inventory	South Korea	An indoor setting at a rehabilitation hospital

Note.

^a We were unable to translate the full text of this study.

^b Only the time relevant for the data analyses was considered.

related to the horticultural intervention. Some participants in Kam and Siu's (2010) study felt fatigued and tired during and after participating in horticultural activities.

3.2. Horticultural interventions compared to other interventions

Five studies found small differences in changes in mean depression scores from baseline to post-intervention between horticultural interventions and other interventions. For example, Makizako et al. (2020) compared their horticultural intervention to a group of people who received classes about traffic safety and disaster prevention (i.e., the educational group) and to a group of people who received an intervention based on physical exercises (i.e., the exercise group). The results from these comparisons were similar, with the horticultural intervention group having a slightly larger reduction in mean depression score from baseline to after the intervention than the two comparison groups ($g = -0.34 [-0.87, 0.20]$, $p = .22$ for the comparison with the educational group, and $g = -0.19 [-0.73, 0.35]$, $p = .49$ for the comparison with the exercise group). In a previous study, Lai et al. (2018) compared their horticultural intervention to a similar intervention (i.e., group size, intervention length, frequency, and duration) involving social activities without using living plants. The authors reported a non-statistically significant difference between the two groups' mean reduction in depressive symptoms ($-0.25 [-1.12, 0.63]$, $p > .05$), and the direction of this effect is unclear. Similar results were found by Vujcic et al. (2017). The authors compared a horticultural intervention to art therapy plus usual care. The authors reported a non-statistically significant difference in mean reduction of depressive symptoms, from pre to post-intervention, between the two groups ($\eta^2 = 0.04$, $p = .31$); again, the direction of this effect is unclear. In another study, Detweiler et al. (2015) compared their horticultural intervention to other occupational activities (e.g., ceramic painting and assembling of leather belts or models in plastic or wood). The horticulture group held a smaller mean depression score after the intervention but the difference with the comparison group was not statistically significant (effect size = .37, $p = .13$). Finally, Kotozaki et al. (2015) compared their horticultural intervention to the provision of stress management sessions and found that the horticulture group held a slightly smaller mean depression score after the intervention ($g = -0.11 [-0.64, 0.42]$, $p = .69$) compared to the alternative intervention group.

Differences in dropout rates between horticultural interventions and other types of interventions were also small, and none of these five studies reported any adverse events related to horticultural interventions. In Makizako et al.'s (2020) study, a few more participants dropped out from the horticultural intervention as compared to the educational group (Risk Ratio = 3.87 [0.46, 32.57], $p = .21$) and to the exercise group (Risk Ratio = 1.33 [0.33, 5.45], $p = .69$). Similarly, Lai et al. (2018) reported that a few more participants dropped out of the horticultural intervention as compared to the group in the

non-horticultural intervention (Risk Ratio = 2.46 [0.50, 12.13], $p = .27$), and Detweiler et al. (2015) found that nine participants dropped out in the comparison group and eight in the horticulture group (Risk Ratio = 0.80 [0.39, 1.62], $p = .54$). In Kotozaki et al. (2015), there were no dropouts.

4. Discussion

In this study, we report evidence from 20 RCTs that assessed the effect of horticultural interventions on adults' depressive symptoms. Unfortunately, we were unable to identify any eligible RCT conducted with children or adolescents through our search strategy. Findings suggest that some horticultural interventions plus usual care may, on average, reduce adults' depressive symptoms more than usual care alone (Fig. 3a). Thirteen of the 15 RCTs assessing this comparison suggested that the addition of horticultural activities to participants' normal daily routines may promote a reduction in their depressive symptoms, and most studies found a moderate or large effect. Two of the 15 RCTs found non-statistically significant differences, on average, in the depressive symptoms of the participants who engaged in horticultural activities and in those who continued their normal routines. These findings are in line with a recent meta-analysis indicating that forest therapy plus usual care may reduce adults' depressive symptoms more than usual care alone (Rosa et al., 2021). Moreover, both studies suggest that people may adhere well to these nature-based interventions (i.e., low dropout rates) and that adverse events are rare.

Several mechanisms could explain why some horticultural interventions reduce adults' depressive symptoms. ART (Kaplan, 1995) and SRT (Ulrich et al., 1991) articulate that positive experiences with nature may reduce people's stress and anxiety and improve mood and attention, all of which are closely related to depression (Fried, 2017; Slavich & Irwin, 2014). For example, higher levels of stress and anxiety have been associated with stronger depressive symptoms (Slavich & Irwin, 2014), and sad mood and concentration problems are both symptoms of major depression (American Psychiatric Association, 2014). Thus, horticulture may reduce people's depressive symptoms by reducing stress and anxiety, and improving mood and concentration. The opportunity to restore one's psychological resources might be linked to the fact that some horticultural interventions are organized outdoors, and likely conducted when weather conditions are favorable. Bad weather conditions limit people's outdoor time, constrain restorative activities, and are linked to more frequent use of antidepressants (Hartig et al., 2007). Related to this, research suggests that sun exposure during nature-based activities may reduce depressive symptoms by improving sleep (Lopresti, 2019; Moreton et al., 2021). Horticultural interventions promote exposure to biodiversity and increase accessibility to plant-based diets, which are also associated with positive health outcomes (Aerts et al., 2018; Leri et al., 2020; Marselle et al., 2021).

Horticultural intervention also involves the practice of physical

Table 3

Description of horticultural activities, comparison group activities, and co-interventions of the randomized controlled trials (RCTs) included in the systematic review.

First author (year)	Horticultural interventions ^a and comparison group activities	Co-interventions	Intervention length in weeks ^b	Intervention frequency ^c	Session duration in hours ^d	Group N
Buru et al. (2021)	Horticultural intervention: Specific gardening activities such as sowing, potting, and planting Usual care: No intervention	No co-intervention was reported	2	Daily	4	8
		Not applicable	Not applicable	Not applicable	Not applicable	8
Pálsdóttir et al. (2020)	Horticultural intervention: Horticulture activities Usual care for stroke survivors	Physical activities and enjoying the garden	10	Two days a week	3.5	48
		Not applicable	Not applicable	Not applicable	Not applicable	44
Kim et al. (2020a)	Horticultural intervention: Activities included transplanting, making bouquets, and harvesting. Usual care for elderly living in a homeless living facility	Walking at the arboretum, reflecting on what changed after the program, and setting goals to live an active and planned life	16	Weekly	1 to 1.5	6
		Not applicable	Not applicable	Not applicable	Not applicable	6
Kim et al. (2020b)	Horticultural intervention: Activities included sowing flower seeds, making a terrarium, and making a scandiamoss tree Usual care: No intervention	Conversations about dementia and therapeutic activities	4	Twice a week	1.5 to 2	10
		Not applicable	Not applicable	Not applicable	Not applicable	9
Makizako et al. (2020)	Horticultural intervention: The program included crop-related activities such as cultivating, growing, and harvesting. Educational group: The classes included topics such as traffic safety and disaster prevention that experts considered less likely to influence study outcomes Exercise group: Each session began with a warm-up period with stretching exercises followed by muscle strength exercises and postural balance re-training.	No co-intervention was reported	20	Weekly	1 to 1.5	26
		Not applicable	26.1	Two times	1.5	28
		Not applicable	20	Weekly	1.5	27
Chu et al. (2019)	Horticultural intervention: Activities included planting seeds, watering plants, and decorating with flowers. Usual care for older residents of nursing homes	A co-intervention was reported but we do not believe it has the potential to reduce participants' depressive symptoms.	8	Weekly	1.5 to 2	75
		Not applicable	Not applicable	Not applicable	Not applicable	75
Najjar et al. (2018)	Horticultural intervention: Activities included planting, watering, and weeding. Usual care for chronically depressed male outpatients	A co-intervention was reported but we do not believe it has the potential to reduce participants' depressive symptoms.	5	Twice a week	2	15
		Not applicable	Not applicable	Not applicable	Not applicable	15
Kim (2018)	Horticultural intervention: The intervention included planting, making crafts with plants, and flower arrangements Usual care: No intervention	No co-intervention was reported	6	Twice a week	1	18
		Not applicable	Not applicable	Not applicable	Not applicable	18
Lai et al. (2018)	Horticultural intervention: The intervention included fertilizing, re-potting plants, watering, trimming, propagation, species introduction, and seeding. Social activities group: All aspects of this group were equivalent to the horticulture group except for the use of living plants.	No co-intervention was reported	8	Weekly	1	46
		Not applicable	8	Weekly	1	50
Ng et al. (2018)	Horticultural intervention: The intervention included gardening, growing, maintaining, and harvesting vegetables and herbs Usual care: No intervention	Guided walking in various parks	26.1	Weekly during 13 weeks then monthly	1	29
		Not applicable	Not applicable	Not applicable	Not applicable	30
Vujcic et al. (2017)	Horticultural intervention: The intervention included plot weeding, potting collecting autumn fruits, and working with plants. Art therapy plus usual care: The control group was included in the occupational and art therapy while continuing to receive conventional therapy, in conditions without plants.	Other activities in contact with nature such as meditation, social support group, and art therapy.	4	Three days a week	1	16
		Not applicable	4	Three days a week	1	14
Kim et al. (2016)	Horticultural intervention: Planting rattan or other plants and creating flower-based decorations	Exercise therapy, cognitive occupational therapy, recollection therapy, art therapy, music therapy, and pharmacological treatment.	26.1	Five times a week	1	32

(continued on next page)

Table 3 (continued)

First author (year)	Horticultural interventions ^a and comparison group activities	Co-interventions	Intervention length in weeks ^b	Intervention frequency ^c	Session duration in hours ^d	Group N
	Usual care for patients with Alzheimer	Not applicable	Not applicable	Not applicable	Not applicable	21
Detweiler et al. (2015)	Horticultural intervention: The intervention included adding soil to garden boxes; planning the types of seeds to plant (e.g., flowers, vegetables, and herbs); planting the seeds; and watering, weeding, and harvesting the vegetables and flowers. Other occupational activities: The group was able to choose from a large variety of crafts, such as ceramic painting, flower arranging, and assembling leather belts or models in plastic or wood.	No co-intervention was reported	3	Five days per week	1	12
		Not applicable	Unclear	Unclear	Unclear	9
Ghanbari et al. (2015)	Horticultural intervention: Plowing land, planting, picking up, and harvesting. Usual care: No intervention	No co-intervention was reported	8.7	Three days a week	1	25
		Not applicable	Not applicable	Not applicable	Not applicable	25
Kotozaki et al. (2015)	Horticultural intervention: The intervention included planting, seeding, watering, weeding, and picking flowers Stress management sessions: These consisted of video lectures regarding stress education	Introductory psychology and stress management lessons	8	Weekly	1	27
		Not applicable	8	Weekly	1	27
Kotozaki (2014)	Horticultural intervention: The intervention included designing a garden planter, seeding, watering, weeding, and picking flowers. Usual care for women victims of an earthquake	No co-intervention was reported	16	Weekly	2	22
		Not applicable	Not applicable	Not applicable	Not applicable	23
Moshfeghi et al. (2014) ^e	Horticultural intervention: Planting, maintaining, and harvesting fruits and vegetables Control group	Unclear whether any co-intervention was reported because we were unable to translate the full text to another language. Not applicable	13	Unclear	Unclear	Unclear
		Not applicable	Not applicable	Not applicable	Not applicable	Unclear
Tse and Ho (2013)	Horticultural intervention: Each participant was responsible for his or her planting, while the research team facilitated and discussed the proper care of the plant, preparing the soils, watering, and adding fertilizers. Usual care for older persons living in a nursing home	Physiotherapy	8	Not reported	Not reported	48
		Not applicable	Not applicable	Not applicable	Not applicable	42
Kam and Siu (2010)	Horticultural intervention: The intervention included watering, fertilizing plants, weeds removal, and loosening soil. Usual care: Participants were receiving workshop training that included a garden tour, and sharing experiences about coping with life events and stress.	No co-intervention was reported	2	Daily	1	10
		Not applicable	Unclear	Unclear	Unclear	12
Kim et al. (2003)	Horticultural intervention: The intervention included planting, transplanting, making flower baskets, and cutting herbs. Usual care for poststroke hemiplegic patients	No co-intervention was reported	6	Five times a week	1	21
		Not applicable	Not applicable	Not applicable	Not applicable	21

Note.

^a Unhealthy participants probably continued their usual treatment while participating in the horticultural interventions.

^b Intervention length refers to the duration of the full intervention.

^c Intervention frequency refers to the frequency of the horticultural activities or comparison group activities.

^d Session duration refers to the duration of the horticultural activities or comparison group activities provided during each session.

^e We were unable to translate the full text of this study.

activity and (typically) some form of socialization, which are both linked to reductions in depressive symptoms (Chu et al., 2019; Clatworthy et al., 2013; Ng et al., 2018; Soga et al., 2017). In line with this, some RCTs that compared engagement in horticultural interventions to physical exercise (Makizako et al., 2020) or to getting involved in social activities (Lai et al., 2018) found small and imprecise (i.e., confidence intervals overlapping zero) differences between these interventions, in terms of reduction in depressive symptoms. This suggests that horticultural interventions are one of several effective, and potentially complementary approaches (e.g., physical activity and socialization), to

improve adults' depressive symptoms. In fact, horticultural interventions were not found to be largely superior to engagement in other occupational activities (Detweiler et al., 2015), art therapy (Vujcic et al., 2017), or stress management sessions (Kotozaki et al., 2015).

4.1. Study limitations

Our findings should be considered in light of several limitations. First, all except one RCT included in our systematic review presented design limitations that might have biased their results (Fig. 2). One

	Risk of bias domains					Overall
	D1	D2	D3	D4	D5	
Buru 2021	-	-	X	X	-	X
Palsdottir 2020	+	+	+	X	+	X
Kim 2020a	-	+	+	X	-	X
Kim 2020b	-	X	X	X	-	X
Makizako 2020	+	+	+	+	+	+
Chu 2019	-	+	+	X	-	X
Najjar 2018	-	X	X	X	-	X
Kim 2018	-	+	+	X	-	X
Lai 2018	+	+	X	+	-	X
Ng 2018	-	-	+	X	+	X
Vujcic 2017	-	X	X	+	-	X
Kim 2016	-	-	X	X	-	X
Detweiler 2015	-	+	X	+	-	X
Ghanbari 2015	X	+	+	X	-	X
Kotozaki 2015	-	-	+	+	X	X
Kotozaki 2014	-	-	+	X	-	X
Tse 2013	-	X	X	X	-	X
Kam 2010	+	+	+	X	-	X
Kim 2003	X	+	+	X	-	X

Domains:
D1: Bias arising from the randomization process.
D2: Bias due to deviations from intended intervention.
D3: Bias due to missing outcome data.
D4: Bias in measurement of the outcome.
D5: Bias in selection of the reported result.

Judgement
 High
 Some concerns
 Low

Fig. 2. Risk of bias of the 19 randomized controlled trials that provided enough data for risk of bias assessment.

limitation presented in all included studies was the inability of keeping the participants unaware of the intervention they were receiving (i.e., blinding). In other words, participants knew when they were receiving the horticultural intervention. This lack of blinding might influence adults' decision to search for additional care if they are not satisfied with the group they were allocated to, or it might bias their reporting of depressive symptoms (Rosa & Delabrida, 2021; Sterne et al., 2016, 2019). Another limitation of many of the included studies was the lack of a registered analysis plan matching the analyses performed in the paper, which would ensure that reporting of results was not selective. Some RCTs did not report enough information to prove that the strategy used to allocate participants to groups was random and concealed (see Rosa, Chaves, Collado, & Harper, 2023; Sterne et al., 2019). Additionally, some RCTs had a considerable amount of missing data from baseline to post-intervention, which can bias the interpretation of an intervention's effect on depressive symptoms under some conditions (Sterne et al., 2016, 2019).

Whereas most included studies may have been affected by some kind of bias, it is unknown how much those biases explain the estimates of horticultural intervention effects that we observed. When considering RCTs with a similar risk of bias, researchers may have more confidence in the efficacy of interventions reported in studies with larger samples and larger estimates than in studies with fewer participants and smaller estimates (Higgins et al., 2019). On average, RCTs included in our review involved about 54 participants, with sample sizes ranging from 12 to 150. To illustrate, both Buru et al. (2021) and Chu et al. (2019) have a high risk of bias, but the latter study included many more participants ($N = 150$) than the first one ($N = 16$) and found a larger estimate of effect ($g = -15.21$ vs. -1.06). Thus, one can be more confident about the efficacy of the intervention reported by Chu et al. (2019) than the one reported by Buru et al. (2021). It is also relevant to note that while a high risk of bias occurs due to limitations in study design, it does not always imply biased estimates (Moustgaard et al., 2020). Future research is essential to understand how study design may influence




Fig. 3. (a) Comparison of the post-intervention mean score or mean change from baseline of horticulture groups versus usual care only using the inverse variance fixed-effect meta-analysis. (b) Comparison of the risk of response to treatment (i.e. ≥50% reduction in depressive symptoms) between horticulture groups and usual care groups, using the Mantel-Haenszel fixed-effect meta-analysis. Events refer to the number of participants who responded to treatment. Green squares refer to standardized mean differences and blue squares to risk ratios. Bigger squares indicated more participants in a study or more events and a bigger diamond indicates greater uncertainty in the combined estimate.

results. Additionally, adherence to relevant Consolidated Standards of Reporting Trials (CONSORT) would improve the interpretation of the results for horticultural intervention studies (Moher et al., 2010).

In addition to these limitations, a systematic review comprises many decisions that influence the interpretation of findings (Higgins et al., 2019). Here we point out how some of our decisions impact the findings' interpretation. First, we did not limit this review to specific populations (e.g., older adults), intervention characteristics (e.g., weekly sessions), and comparison groups (e.g., usual care). Hence, the included RCTs are different in important characteristics that somewhat preclude a comprehensive quantitative synthesis of all results (i.e., meta-analysis). We, therefore, chose to present a forest plot with effect estimates from the RCTs comparing horticultural interventions plus usual care with just usual care (Fig. 3a). Nonetheless, we recommend that readers do not focus on the combined estimate from these studies. Instead, they may consider how different kinds of horticultural interventions (including the kind of activities provided, their length, frequency, and duration) may improve the depressive symptoms of specific groups (e.g., older adults) as compared to the alternative interventions (i.e., usual care). More randomized studies that isolate the impacts of specific variables are needed to improve the understanding of how participants' and interventions' characteristics may influence the study results.

Also linked to our broad criteria of eligibility, we included studies

independent of whether or not their participants had a diagnosis of depression. We did this because every individual can experience depressive symptoms (e.g., sad mood) to a certain degree. Some studies included participants diagnosed with mental health problems related to but not necessarily involving just depression, such as adults with psychiatric illnesses (Kam & Siu, 2010; Vujcic et al., 2017). In fact, only one (Najjar et al., 2018) out of the 20 RCTs included exclusively adults diagnosed with depression. Thus, more RCTs with individuals exclusively diagnosed with depression are needed.

Finally, concerning our methodology, one researcher conducted the title and abstract screening. This approach was efficient, but the risk of unintentionally excluding a potentially relevant study might have been reduced if two researchers were involved in this process. Unfortunately, this was a necessary decision to facilitate the execution of this systematic review. Additionally, no systematic review is expected to include all studies relevant to the research question since no search strategy is perfectly effective (Higgins et al., 2019).

5. Conclusion and next steps

To date, our systematic review is the most comprehensive summary of studies estimating the effect of horticultural interventions on adults' depressive symptoms. We found relatively consistent results indicating

Table 4

Percentage of change from baseline in depression scores, standardized mean change, number of participants who had a $\geq 50\%$ reduction on depression scores from baseline to post-intervention (i.e., responders), and dropouts in the horticultural interventions and comparison groups of the randomized controlled trials (RCTs) included in this systematic review.

First author (year)	Group	Percentage of change from baseline ^a	Standardized mean change ^b	Responders ^c	Dropouts
Buru et al. (2021)	Horticultural intervention	-43.8	-1.63	3/8	Unclear
	Usual care	-10.8	-1.18	2/8	18/32
Pálsdóttir et al. (2020)	Horticultural intervention	-19.4	NR	NR	1/51
	Usual care	-20.1	NR	NR	7/50
Kim et al. (2020a)	Horticultural intervention	-17.2	-0.32	NR	0/6
	Usual care	7.3	0.13	NR	0/6
Kim et al. (2020b)	Horticultural intervention	-6.8	-0.22	NR	Unclear
	Usual care	11.1	0.51	NR	Unclear
Makizako et al. (2020)	Horticultural intervention	-31.9	-0.47	NR	4/30
	Exercise group	-25.4	-0.72	NR	3/30
	Educational group	-20.3	-0.52	NR	1/29
Chu et al. (2019)	Horticultural intervention	-62.9	-12.43	NR	0/75
	Usual care	48.5	6.95	NR	0/75
Najjar et al. (2018)	Horticultural intervention	-25.2	-1.05	NR	0/15
	Usual care	1.7	0.06	NR	0/15
Kim (2018)	Horticultural intervention	-25.3	-1.31	NR	0/18
	Usual care	0.7	0.69	NR	0/18
Lai et al. (2018)	Horticultural intervention	NR	NR	NR	5/56
	Social activities	NR	NR	NR	2/55
Ng et al. (2018)	Horticultural intervention	NR	NR	NR	0/29
	Usual care	NR	NR	NR	0/30
Vujcic et al. (2017)	Horticultural intervention	NR	NR	NR	NR
	Art therapy plus usual care	NR	NR	NR	NR
Kim et al. (2016)	Horticultural intervention	-8.8	-0.19	NR	0/32
	Usual care	-0.7	-0.02	NR	Unclear
Detweiler et al. (2015)	Horticultural intervention	NR	NR	NR	8/20
	Other occupational activities	NR	NR	NR	9/18
Ghanbari et al. (2015)	Horticultural intervention	-51.5	-1.35	13/25	0/25
	Usual care	-13.7	-0.58	0/25	0/25
Kotozaki et al. (2015)	Horticultural intervention	-12.1	-0.23	NR	0/27
	Stress management sessions	-14.0	-0.30	NR	0/27
Kotozaki (2014)	Horticultural intervention	-41.9	-0.69	NR	0/22
	Usual care	-15.2	-0.22	NR	0/23
Moshfeghi et al. (2014) ^d	Horticultural intervention	Unclear	Unclear	Unclear	Unclear
	Control group	Unclear	Unclear	Unclear	Unclear
Tse and Ho (2013)	Horticultural intervention	-29.6	-0.63	NR	NR
	Usual care	7.0	0.12	NR	NR
Kam and Siu (2010)	Horticultural intervention	-63.0	-1.01	NR	2/12
	Usual care	-12.6	-0.13	NR	0/12
Kim et al. (2003)	Horticultural intervention	-62.7	-4.89	19/21	0/21
	Usual care	-58.8	-3.57	15/21	0/21

Note. Negative values for change from baseline and standardized mean change signify reductions in depressive symptoms.

^a Change in score divided by baseline score times 100.

^b Change in score divided by the baseline standard deviation.

^c Having a $\geq 50\%$ reduction in depressive symptoms from baseline to post-intervention; estimated using the formulae described by Furukawa et al. (2005).

^d We were unable to translate the full text of this study. NR = Not reported.

that horticultural interventions plus usual care may reduce adults' depressive symptoms more than usual care alone. Overall, we observed some variability in the magnitude of the effect estimates across the included RCTs, which might be due to variability in participants, interventions, and the outcome measures used. We were unable to determine which specific characteristics of the participants, interventions, or outcome measures are associated with a stronger impact of horticultural interventions on depressive symptoms. Thus, we encourage researchers to conduct RCTs aimed at exploring the potential influence these characteristics have on the effect of horticultural interventions on depression. A randomized study could provide a similar intervention to two different groups of individuals or a slightly different intervention to the same participants. For instance, future RCTs could assess the relevance of sun exposure for improvement in depressive symptoms during horticultural interventions by comparing groups randomly allocated to indoor versus outdoor settings. As another example, future RCTs could compare whether group-based horticultural interventions are more effective than participation in one-on-one, or solo horticultural activities. Such an investigation could provide extra support to the evidence that social interactions play a role in reducing

depressive symptoms during horticultural interventions (Chu et al., 2019; Clatworthy et al., 2013; Lin et al., 2020; Ng et al., 2018; Soga et al., 2017).

Our findings also suggest that people may adhere well to horticultural interventions (i.e., low dropout rates) and that adverse events like fatigue and tiredness (Kam & Siu, 2010) during and after these interventions are likely rare. Nonetheless, we highlight that other complementary interventions, such as the practice of physical exercise (Makizako et al., 2020) and social activities without direct interaction with plants (Lai et al., 2018), might provide similar, but maybe slightly smaller reductions in adults' depressive symptoms. Given the design limitations of virtually all studies, more rigorous RCTs are needed. It may be worth conducting RCTs in places where the effect of horticultural interventions has been scarcely examined, like Latin America, Africa, and Oceania. It may also be prudent to focus RCTs on people diagnosed with depression, as well as young people, especially because we did not find any RCTs (eligible for our systematic review) involving children or adolescents.

Future systematic reviews could also explore other outcomes relevant to understanding the potential value of horticultural interventions,

including the possible effects of these activities on other mental (e.g., anxiety, loneliness, and anger) and physical outcomes (e.g., weight loss). Systematic reviews that directly assess the effect of horticultural interventions on specific symptoms of depression (e.g., sad mood and anhedonia) are also warranted since our review focused on aggregate scores from depression outcome measures, not on specific symptoms. Finally, studies should consider the financial cost, and relative benefits, of implementing horticultural interventions compared to other more conventional strategies commonly employed to prevent or treat depression and other mental health disorders.

Funding

This systematic review was registered in the Open Science Framework (<https://archive.org/details/osf-registrations-tyk7m-v1>). We have no conflict of interest to disclose. This study was partially funded by the Spanish Ministry of Science, Innovation, and Universities (PGC 2018-095502-B-I00) and by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

Declaration of competing interest

The authors declare they have no conflict of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jenvp.2023.102112>.

References

- Aerts, R., Honnay, O., & Van Nieuwenhuysse, A. (2018). Biodiversity and human health: Mechanisms and evidence of the positive health effects of diversity in nature and green spaces. *British Medical Bulletin*, 127(1), 5–22. <https://doi.org/10.1093/bmb/ldy021>
- American Horticultural Therapy Association. (2017). *AHTA definitions and positions paper*.
- American Psychiatric Association. (2014). In *Manual diagnóstico e estatístico de transtornos mentais: DSM-5 [diagnostic and statistical manual of mental disorders: DSM-5]* (5th ed.). Artmed.
- Berk, M., & Jacka, F. N. (2019). Diet and depression—from confirmation to implementation. *JAMA*, 321(9), 842–843. <https://doi.org/10.1001/jama.2019.0273>
- Berto, R. (2014). The role of nature in coping with psycho-physiological stress: A literature review on restorativeness. *Behavioral Sciences*, 4(4), 394–409. <https://doi.org/10.3390/bs4040394>
- Buru, T., Kállay, É., Olar, L. E., Ștefan, R., Cantor, M., Papuc, I., & Buta, E. (2021). Studies regarding the influence of therapeutic horticulture on the human-nature relationship and the increase of well-being. *Acta Horticulturae*, 1330, 75–86. <https://doi.org/10.17660/ActaHortic.2021.1330.10>
- Catalan-Matamoros, D., Gomez-Conesa, A., Stubbs, B., & Vancampfort, D. (2016). Exercise improves depressive symptoms in older adults: An umbrella review of systematic reviews and meta-analyses. *Psychiatry Research*, 244, 202–209. <https://doi.org/10.1016/j.psychres.2016.07.028>
- Chu, H.-Y., Chen, M.-F., Tsai, C.-C., Chan, H.-S., & Wu, T.-L. (2019). Efficacy of a horticultural activity program for reducing depression and loneliness in older residents of nursing homes in Taiwan. *Geriatric Nursing*, 40(4), 386–391. <https://doi.org/10.1016/j.gerinurse.2018.12.012>
- Cipriani, J., Benz, A., Holmgren, A., Kinter, D., McGarry, J., & Rufino, G. (2017). A systematic review of the effects of horticultural therapy on persons with mental health conditions. *Occupational Therapy in Mental Health*, 33(1), 47–69. <https://doi.org/10.1080/0164212X.2016.1231602>
- Cipriani, A., Furukawa, T. A., Salanti, G., Chaimani, A., Atkinson, L. Z., Ogawa, Y., Leucht, S., Ruhe, H. G., Turner, E. H., Higgins, J. P. T., Egger, M., Takeshima, N., Hayasaka, Y., Imai, H., Shinohara, K., Tajika, A., Ioannidis, J. P. A., & Geddes, J. R. (2018). Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *The Lancet*, 391(10128), 1357–1366. [https://doi.org/10.1016/S0140-6736\(17\)32802-7](https://doi.org/10.1016/S0140-6736(17)32802-7)
- Clatworthy, J., Hinds, J., & Camic, P. M. (2013). Gardening as a mental health intervention: A review. *Mental Health Review Journal*, 18(4), 214–225. <https://doi.org/10.1108/MHRJ-02-2013-0007>
- Crossan, C., & Salmoni, A. (2021). A simulated walk in nature: Testing predictions from the attention restoration theory. *Environment and Behavior*, 53(3), 277–295. <https://doi.org/10.1177/0013916519882775>
- Cuijpers, P., Noma, H., Karyotaki, E., Vinkers, C. H., Cipriani, A., & Furukawa, T. A. (2020). A network meta-analysis of the effects of psychotherapies, pharmacotherapies and their combination in the treatment of adult depression. *World Psychiatry*, 19(1), 92–107. <https://doi.org/10.1002/wps.20701>
- Detweiler, M. B., Self, J. A., Lane, S., Spencer, L., Lutgens, B., Kim, D.-Y., Halling, M. H., Rudder, T. F., & Lehmann, L. (2015). Horticultural therapy: A pilot study on modulating cortisol levels and indices of substance craving, posttraumatic stress disorder, depression, and quality of life in veterans. *Alternative Therapies in Health & Medicine*, 21(4), 36–41.
- Ferne, C. R., Gabrielsen, L. E., Andersen, A. J. W., & Mesel, T. (2017). Unpacking the black box of wilderness therapy. *Qualitative Health Research*, 27(1), 114–129. <https://doi.org/10.1177/1049732316655776>
- Fried, E. I. (2017). The 52 symptoms of major depression: Lack of content overlap among seven common depression scales. *Journal of Affective Disorders*, 208, 191–197. <https://doi.org/10.1016/j.jad.2016.10.019>
- Frost, S., Kannis-Dymand, L., Schaffer, V., Milleer, P., Allen, A., Stallman, H., Mason, J., Wood, A., & Atkinson-Nolte, J. (2022). Virtual immersion in nature and psychological well-being: A systematic literature review. *Journal of Environmental Psychology*, 80(101765), 1–9. <https://doi.org/10.1016/j.jenvp.2022.101765>
- Furukawa, T. A., Cipriani, A., Barbui, C., Brambilla, P., & Watanabe, N. (2005). Imputing response rates from means and standard deviations in meta-analyses. *International Clinical Psychopharmacology*, 20(1), 49–52. <https://doi.org/10.1097/00004850-200501000-00010>
- Ghanbari, S., Jafari, F., Bagheri, N., Neamtollahi, S., & Shayanpour, R. (2015). Study of the effect of using purposeful activity (gardening) on depression of female resident in Golestan Dormitory of Ahvaz Jundishapur University of Medical Sciences. *Journal of Rehabilitation Sciences & Research*, 2(1), 8–11. <https://doi.org/10.30476/JRSR.2015.41066>
- Hartig, T. (2021). Restoration in nature: Beyond the conventional narrative. In A. R. Schutte, J. Torquati, & J. R. Stevens (Eds.), *Nature and psychology: Biological, cognitive, developmental, and social pathways to well-being*. Springer.
- Hartig, T., Catalano, R., & Ong, M. (2007). Cold summer weather, constrained restoration, and the use of antidepressants in Sweden. *Journal of Environmental Psychology*, 27(2), 107–116. <https://doi.org/10.1016/j.jenvp.2007.02.002>
- Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., & Welch, V. A. (Eds.). (2019). *Cochrane handbook for systematic reviews of interventions* (2nd ed.). John Wiley & Sons.
- Houge Mackenzie, S., Hodge, K., & Filep, S. (2021). How does adventure sport tourism enhance well-being? A conceptual model. *Tourism Recreation Research*, 1–14. <https://doi.org/10.1080/02508281.2021.1894043>
- Jiang, B., He, J., Chen, J., Larsen, L., & Wang, H. (2021). Perceived green at speed: A simulated driving experiment raises new questions for attention restoration theory and stress reduction theory. *Environment and Behavior*, 53(3), 296–335. <https://doi.org/10.1177/0013916520947111>
- Kamioka, H., Tsutani, K., Yamada, M., Park, H., Okuizumi, H., Honda, T., Okada, S., Park, S.-J., Kitayuguchi, J., Abe, T., Handa, S., & Mutoh, Y. (2014). Effectiveness of horticultural therapy: A systematic review of randomized controlled trials. *Complementary Therapies in Medicine*, 22(5), 930–943. <https://doi.org/10.1016/j.ctim.2014.08.009>
- Kam, M. C. Y., & Siu, A. M. H. (2010). Evaluation of a horticultural activity programme for persons with psychiatric illness. *Hong Kong Journal of Occupational Therapy*, 20(2), 80–86. [https://doi.org/10.1016/S1569-1861\(11\)70007-9](https://doi.org/10.1016/S1569-1861(11)70007-9)
- Kaplan, S. (1995). The restorative benefits of nature: Toward an integrative framework. *Journal of Environmental Psychology*, 15(3), 169–182. [https://doi.org/10.1016/0272-4944\(95\)90001-2](https://doi.org/10.1016/0272-4944(95)90001-2)
- Kim, S.-Y., Son, K.-C., Jung, H.-J., Yoo, J.-H., Kim, B.-S., & Park, S.-W. (2003). Effect of horticultural therapy on functional rehabilitation in hemiplegic patients after stroke. *Journal of the Korean Society for Horticultural Science*, 44(5), 780–785.
- Kim, W., Lim, S.-K., Chung, E.-J., & Woo, J.-M. (2009). The effect of cognitive behavior therapy-based psychotherapy applied in a forest environment on physiological changes and remission of major depressive disorder. *Psychiatry Investigation*, 6(4), 245. <https://doi.org/10.4306/pi.2009.6.4.245>
- Kim, H.-J., Yang, Y., Oh, J.-G., Oh, S., Choi, H., Kim, K. H., & Kim, S. H. (2016). Effectiveness of a community-based multidomain cognitive intervention program in patients with Alzheimer's disease. *Geriatrics and Gerontology International*, 16(2), 191–199. <https://doi.org/10.1111/ggi.12453>
- Kim, Y. H., Lee, S.-H., Park, C.-S., Bae, H.-O., Kim, Y. J., & Huh, M. R. (2020a). A horticultural therapy program focusing on gardening activities to promote psychological, emotional and social health of the elderly living in a homeless living facility for a long time: A pilot study. *Journal of People, Plants, and Environment*, 23(5), 565–576. <https://doi.org/10.11628/ksppe.2020.23.5.565>
- Kim, Y. H., Park, C. S., Bae, H.-O., Lim, E. J., Kang, K. H., Lee, E. S., ... Huh, M. R. (2020b). Horticultural therapy programs enhancing quality of life and reducing depression and burden for caregivers of elderly with dementia. *Journal of People, Plants, and Environment*, 23(3), 305–320. <https://doi.org/10.11628/ksppe.2020.23.3.305>
- Kotzaki, Y. (2014). Horticultural therapy as a measure for recovery support of regional community in the disaster area: a preliminary experiment for forty five women who living certain region in the coastal area of Miyagi Prefecture. *International Journal of Emergency Mental Health*, 16(2), 284–287.
- Kotzaki, Y., Takeuchi, H., Sekiguchi, A., Araki, T., Takahashi, K., Yamamoto, Y., Nozawa, T., Taki, Y., & Kawashima, R. (2015). Positive effects of the victim by the growing of plants after great east Japan earthquake. *International Journal of Recent Scientific Research*, 6(2), 2850–2858.
- Lai, C. K. Y., Kwan, R. Y. C., Lo, S. K. L., Fung, C. Y. Y., Lau, J. K. H., & Tse, M. M. Y. (2018). Effects of horticulture on frail and prefrail nursing home residents: A randomized controlled trial. *Journal of the American Medical Directors Association*, 19(8), 696–702. <https://doi.org/10.1016/j.jamda.2018.04.002>

- Leri, M., Scuto, M., Ontario, M. L., Calabrese, V., Calabrese, E. J., Bucciantini, M., & Stefani, M. (2020). Healthy effects of plant polyphenols: Molecular mechanisms. *International Journal of Molecular Sciences*, 21(1250), 1–40. <https://doi.org/10.3390/ijms21041250>
- Lin, T.-Y., Huang, C.-M., Hsu, H.-P., Liao, J.-Y., Cheng, V. Y.-W., Wang, S.-W., & Guo, J.-L. (2020). Effects of a combination of three-dimensional virtual reality and hands-on horticultural therapy on institutionalized older adults' physical and mental health: Quasi-experimental design. *Journal of Medical Internet Research*, 22(11), Article e19002. <https://doi.org/10.2196/19002>
- Liu, Y., Li, B., Sampson, S. J., Roberts, S., Zhang, G., & Wu, W. (2014). Horticultural therapy for schizophrenia. *Cochrane Database of Systematic Reviews*, 2019(9). <https://doi.org/10.1002/14651858.CD009413.pub2>
- Lopresti, A. L. (2019). It is time to investigate integrative approaches to enhance treatment outcomes for depression? *Medical Hypotheses*, 126, 82–94. <https://doi.org/10.1016/j.mehy.2019.03.008>
- Makizako, H., Tsutsumimoto, K., Doi, T., Makino, K., Nakakubo, S., Liu-Ambrose, T., & Shimada, H. (2020). Exercise and horticultural programs for older adults with depressive symptoms and memory problems: A randomized controlled trial. *Journal of Clinical Medicine*, 9(1). <https://doi.org/10.3390/jcm9010099>
- Marselle, M. R., Hartig, T., Cox, D. T. C., de Bell, S., Knapp, S., Lindley, S., Triguero-Mas, M., Böhning-Gaese, K., Braubach, M., Cook, P. A., de Vries, S., Heintz-Buschart, A., Hofmann, M., Irvine, K. N., Kabisch, N., Kolek, F., Kraemer, R., Markevych, I., Martens, D., ... Bonn, A. (2021). Pathways linking biodiversity to human health: A conceptual framework. *Environment International*, 150(106420), 1–22. <https://doi.org/10.1016/j.envint.2021.106420>
- McCormack, J., & Korownyk, C. (2018). Effectiveness of antidepressants. *BMJ*, k1073. <https://doi.org/10.1136/bmj.k1073>
- McGuinness, L. (2019). robvis: An R package and web application for visualising risk-of-bias assessments. <https://github.com/mcguinlu/robvis>.
- Moher, D., Hopewell, S., Schulz, K. F., Montori, V., Gøtzsche, P. C., Devereaux, P. J., Elbourne, D., Egger, M., & Altman, D. G. (2010). CONSORT 2010 explanation and elaboration: Updated guidelines for reporting parallel group randomised trials. *International Journal of Surgery*, 10(1), 28–55. <https://doi.org/10.1136/bmj.c869>
- Moll, A., Collado, S., Staats, H., & Corraliza, J. A. (2022). Restorative effects of exposure to nature on children and adolescents: A systematic review. *Journal of Environmental Psychology*, 84, Article 101884. <https://doi.org/10.1016/j.jenvp.2022.101884>
- Moreton, S. G., Brennan, M. K., Nicholls, V. I., Wolf, I. D., & Muir, D. L. (2021). Exploring potential mechanisms underpinning the therapeutic effects of surfing. *Journal of Adventure Education and Outdoor Learning*, 1–18. <https://doi.org/10.1080/14729679.2021.1884104>
- Morris, S. B. (2008). Estimating effect sizes from pretest-posttest-control group designs. *Organizational Research Methods*, 11(2), 364–386. <https://doi.org/10.1177/1094428106291059>
- Moshfeghi, G., Rezaabakhsh, H., & Danesh, E. (2014). Effectiveness of horticulture therapy on depression among older adults in nursing homes. *Scientific-Research Journal of the Faculty of Nursing and Midwifery*, 86.
- Moustgaard, H., Clayton, G. L., Jones, H. E., Boutron, I., Jørgensen, L., Laursen, D. R. T., Olsen, M. F., Paludan-Müller, A., Ravaut, P., Savović, J., Sterne, J. A. C., Higgins, J. P. T., & Hróbjartsson, A. (2020). Impact of blinding on estimated treatment effects in randomised clinical trials: Meta-epidemiological study. *BMJ*, Article 16802. <https://doi.org/10.1136/bmj.16802>
- Murroni, V., Cavalli, R., Basso, A., Borella, E., Meneghetti, C., Melendugno, A., & Pazzaglia, F. (2021). Effectiveness of therapeutic gardens for people with dementia: A systematic review. *International Journal of Environmental Research and Public Health*, 18(18), 9595. <https://doi.org/10.3390/ijerph18189595>
- Najjar, A. H., Foroozandeh, E., & Gharnah, H. A. A. (2018). Horticulture therapy effects on memory and psychological symptoms of depressed male outpatients. *Iranian Rehabilitation Journal*, 16(2), 147–154. <https://doi.org/10.32598/irj.16.2.147>
- Ng, K. S. T., Sia, A., Ng, M. K. W., Tan, C. T. Y., Chan, H. Y., Tan, C. H., Rawtaer, I., Feng, L., Mahendran, R., Larbi, A., Kua, E. H., & Ho, R. C. M. (2018). Effects of horticultural therapy on asian older adults: A randomized controlled trial. *International Journal of Environmental Research and Public Health*, 15(1715). <https://doi.org/10.3390/ijerph15081705>
- Nicholas, S. O., Giang, A. T., & Yap, P. L. K. (2019). The effectiveness of horticultural therapy on older adults: A systematic review. *Journal of the American Medical Directors Association*, 20(10), 1351.e1–1351.e11. <https://doi.org/10.1016/j.jamda.2019.06.021>
- Ohly, H., White, M. P., Wheeler, B. W., Bethel, A., Ukoumunne, O. C., Nikolaou, V., & Garside, R. (2016). Attention restoration theory: A systematic review of the attention restoration potential of exposure to natural environments. *Journal of Toxicology and Environmental Health, Part A*, 19(7), 305–343. <https://doi.org/10.1080/10937404.2016.1196155>
- Owens, M., & Bunce, H. L. I. (2022). The potential for outdoor nature-based interventions in the treatment and prevention of depression. *Frontiers in Psychology*, 13. <https://doi.org/10.3389/fpsyg.2022.740210>
- Pálsdóttir, A. M., Stigmar, K., Norrving, B., Petersson, I. F., Åström, M., & Pessah-Rasmussen, H. (2020). The nature stroke study; NASTRU: A randomized controlled trial of nature-based post-stroke fatigue rehabilitation. *Journal of Rehabilitation Medicine*, 52(2), Article jrm00020. <https://doi.org/10.2340/16501977-2652>
- Reese, R. F., & Gosling, M. (2020). The group EcoWellness model of change: A conceptual framework for facilitating groups in nature. *Journal for Specialists in Group Work*, 45(4), 331–352. <https://doi.org/10.1080/01933922.2020.1799465>
- Review Manager (RevMan) [Computer program] (5.4.1). (2020). The Cochrane Collaboration.
- Riedel, M., Möller, H.-J., Obermeier, M., Schennach-Wolff, R., Bauer, M., Adli, M., Kronmüller, K., Nickel, T., Brieger, P., Laux, G., Bender, W., Heuser, I., Zeiler, J., Gaebel, W., & Seemüller, F. (2010). Response and remission criteria in major depression – a validation of current practice. *Journal of Psychiatric Research*, 44(15), 1063–1068. <https://doi.org/10.1016/j.jpsychires.2010.03.006>
- Rosa, C. D., Chaves, T. S., Collado, S., & Harper, N. J. (2023). Improving the analysis and reporting of studies of nature-based adventure interventions: A review of studies published in JAEOL. *Journal of Adventure Education and Outdoor Learning*, 1–20. <https://doi.org/10.1080/14729679.2023.2196638>
- Rosa, C. D., Chaves, T. S., Collado, S., Larson, L. R., & Profice, C. C. (2023). The effect of nature-based adventure interventions on depression: A systematic review. *Environment and Behavior*, 1–35. <https://doi.org/10.1177/00139165231174615>
- Rosa, C. D., & Delabrida, Z. (2021). Método experimental e ensaios clínicos: Metassíntese de artigos de revisão publicados em português [Experimental method and clinical trials: Metasynthesis of review articles published in Portuguese]. *Psico*, 52(4), Article e36259. <https://doi.org/10.15448/1980-8623.2021.4.36259>
- Rosa, C. D., Larson, L. R., Collado, S., & Profice, C. C. (2021). Forest therapy can prevent and treat depression: Evidence from meta-analyses. *Urban Forestry and Urban Greening*, 57, Article 126943. <https://doi.org/10.1016/j.ufug.2020.126943>
- Russell, K. C., & Farnum, J. (2004). A concurrent model of the wilderness therapy process. *Journal of Adventure Education and Outdoor Learning*, 4(1), 39–55. <https://doi.org/10.1080/14729670485200411>
- Shamseer, L., Moher, D., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., Shekelle, P., & Stewart, L. A. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *BMJ*, 349, g7647. <https://doi.org/10.1136/bmj.g7647>
- Shao, Y., Elsadek, M., & Liu, B. (2020). Horticultural activity: Its contribution to stress recovery and wellbeing for children. *International Journal of Environmental Research and Public Health*, 17(4). <https://doi.org/10.3390/ijerph17041229>
- Shin, W. S., Shin, C. S., & Yeoun, P. S. (2012). The influence of forest therapy camp on depression in alcoholics. *Environmental Health and Preventive Medicine*, 17(1), 73–76. <https://doi.org/10.1007/s12199-011-0215-0>
- Slavich, G. M., & Irwin, M. R. (2014). From stress to inflammation and major depressive disorder: A social signal transduction theory of depression. *Psychological Bulletin*, 140(3), 774–815. <https://doi.org/10.1037/a0035302>
- Soga, M., Gaston, K. J., & Yamaura, Y. (2017). Gardening is beneficial for health: A meta-analysis. *Preventive Medicine Reports*, 5, 92–99. <https://doi.org/10.1016/j.pmedr.2016.11.007>
- Sterne, J. A., Hernán, M. A., Reeves, B. C., Savović, J., Berkman, N. D., Viswanathan, M., Henry, D., Altman, D. G., Ansari, M. T., Boutron, I., Carpenter, J. R., Chan, A.-W., Churchill, R., Deeks, J. J., Hróbjartsson, A., Kirkham, J., Jüni, P., Loke, Y. K., Pigott, T. D., ... Higgins, J. P. (2016). ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*, Article 14919. <https://doi.org/10.1136/bmj.i4919>
- Sterne, J. A., Savović, J., Page, M. J., Elbers, R. G., Blencowe, N. S., Boutron, I., Cates, C. J., Cheng, H.-Y., Corbett, M. S., Eldridge, S. M., Emberson, J. R., Hernán, M. A., Hopewell, S., Hróbjartsson, A., Junqueira, D. R., Jüni, P., Kirkham, J. J., Lasserson, T., Li, T., ... Higgins, J. P. T. (2019). RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ*, Article 14898. <https://doi.org/10.1136/bmj.14898>
- Sturm, J., Plöderl, M., Fartacek, C., Kralovec, K., Neunhäuserer, D., Niederseer, D., Hitzl, W., Niebauer, J., Schiepek, G., & Fartacek, R. (2012). Physical exercise through mountain hiking in high-risk suicide patients. A randomized crossover trial. *Acta Psychiatrica Scandinavica*, 126(6), 467–475. <https://doi.org/10.1111/j.1600-0447.2012.01860.x>
- Tse, M. M. Y., & Ho, S. S. K. (2013). Pain management for older persons living in nursing homes: A pilot study. *Pain Management Nursing*, 14(2). <https://doi.org/10.1016/j.pmn.2011.01.004>
- Tu, H. (2022). Effect of horticultural therapy on mental health: A meta-analysis of randomized controlled trials. *Journal of Psychiatric and Mental Health Nursing*, 29(4), 603–615. <https://doi.org/10.1111/jpm.12818>
- Ulrich, R. S., Simons, R. F., Losito, B. D., Fiorito, E., Miles, M. A., & Zelson, M. (1991). Stress recovery during exposure to natural and urban environments. *Journal of Environmental Psychology*, 11(3), 201–230. [https://doi.org/10.1016/S0272-4944\(05\)80184-7](https://doi.org/10.1016/S0272-4944(05)80184-7)
- Vickers, A. J. (2001). The use of percentage change from baseline as an outcome in a controlled trial is statistically inefficient: A simulation study. *BMC Medical Research Methodology*, 1(1), 6. <https://doi.org/10.1186/1471-2288-1-6>
- Vujcic, M., Tomicevic-Dubljevic, J., Grbic, M., Lecic-Tosevski, D., Vukovic, O., & Toskovic, O. (2017). Nature based solution for improving mental health and well-being in urban areas. *Environmental Research*, 158, 385–392. <https://doi.org/10.1016/j.envres.2017.06.030>
- Wang, D., & MacMillan, T. (2013). The benefits of gardening for older adults: A systematic review of the literature. *Activities, Adaptation & Aging*, 37(2), 153–181. <https://doi.org/10.1080/01924788.2013.784942>
- Wang, Z., Zhang, Y., Lu, S., Tan, L., Guo, W., Lown, M., Hu, X., & Liu, J. (2022). Horticultural therapy for general health in the older adults: A systematic review and meta-analysis. *PLoS One*, 17(2), Article e0263598. <https://doi.org/10.1371/journal.pone.0263598>
- Wilson, E. O. (1984). *Biophilia*. MIT Press.
- World Health Organization. (2017). *Depression and other common mental disorders: Global health estimates*. <https://www.who.int/publications/i/item/depression-global-health-estimates>.
- World Health Organization. (2022). *Mental Health and COVID-19: Early evidence of the pandemic's impact*. World Health Organization.